



Retrospective evaluation of correlation and agreement between two recovery scoring systems in horses

Stefania Scarabelli, Eva Rioja

Recovery is one of the most important and risky phases in equine anaesthesia.¹ Several recovery scoring systems (RSS) have been developed to evaluate risk factors associated with morbidity and mortality during this perioperative phase. A descriptive scale² and a composite rating scale³ are commonly used in equine practice and have been reported as reliable methods for qualifying recovery from general anaesthesia in horses⁴; however, a truly validated and universally used RSS is not available at present, complicating the comparison between different studies in which different scales are used.

The aim of this study was to add information to the existing knowledge about recovery in horses, evaluating correlation and agreement between the two RSS used at the authors' institution.

Anaesthetic records of horses undergoing general anaesthesia between January 2013 and November 2015 were retrospectively reviewed. During this period, two types of recovery scores were used: Y, modified from Young and Taylor,² and D, modified from Donaldson and others.³ System Y (Table 1) is a simple descriptive scale that employs a scale from 1 to 5 to score recovery quality in its entirety; it assigns several descriptors to each score, and high numerical values indicate 'worst' recovery.

System D (Table 2) is a composite scoring system in which eight phases of the recovery are scored and then the values are summated to obtain a total score that is then matched to an overall descriptive score developed to rank the recovery on a scale from 1 to 5 according to

the quality of recovery, with the highest number of the descriptive score indicating the worst recovery, similar to system Y.⁵

All horses included in the study had unassisted recoveries. For each horse, recoveries were scored with both systems by the same person, whose identity was not specified upon the anaesthetic record but that was either a diplomate of the European or American College of Veterinary Anaesthesia and Analgesia, a resident in Veterinary Anaesthesia and Analgesia, or a final-year student.

Horses were included in the study if data about American Society of Anaesthesiologists (ASA) grade,⁶ type of surgery (emergency or elective) and both recovery scores were reported in the anaesthetic records. Demographic data of the horses were also recorded.

Data were analysed using commercially available statistical software (SPSS, V.22.0, 2013). Correlation and agreement were evaluated with Spearman's coefficient and *k*-weighted, respectively. These analyses were performed in all the cases grouped together and after subdividing the horses into groups based on ASA category and type of procedure.

TABLE 1: System Y (modified from Young and Taylor²)

Score 1	Excellent: perfect, no ataxia; stands at first attempt as if rising from rest
Score 2	Good: stands after first or second attempt, some weakness/ataxia and may knuckle slightly after standing but not in danger of falling over
Score 3	Fair: stands after several attempts, crashes/ricochets around recovery box in excited/delirious manner often giving moments for concerns; moderate weakness/ataxia and may knuckle on fetlocks after standing; looks very wobbly and may teeter occasionally, but if falls, gets up again almost immediately; may sustain minor cuts/grazes but not requiring serious treatment intervention (not requiring suturing or further general anaesthesia to treat)
Score 4	Marginal: multiple attempts to stand, crashes/ricochets around recovery box in excited, almost manic manner giving rise to concern; very ataxic when finally standing and commonly falls down again; sustains injury requiring intervention, and which may compromise horse's future/use
Score 5	Poor/unsuccessful: multiple attempts to stand, crashes/ricochets around recovery box in excited, manic manner perhaps until exhausted; may have multiple frantic attempts to stand between exhausted episodes; fails to stand because of postanaesthetic myopathy/neuropathy or through sustaining major/terminal injury (eg, long bone fracture, neck or skull-base fracture, dislocation)

Veterinary Record (2018)

doi: 10.1136/vr.104546

School of Veterinary Science, Philip Leverhulme Equine Hospital, University of Liverpool, Neston, UK

E-mail for correspondence: Liverpool University; scara@liverpool.ac.uk

SS and ER contributed equally.

Provenance and peer review Not commissioned; externally peer reviewed.

Received June 2, 2017
Revised September 20, 2017
Accepted October 29, 2017

TABLE 2: System D (modified from Donaldson and others and Valverde and others^{3,5})

Overall attitude	1-Calm 2-Calm/determined 3-Confused, dizzy 4-Frantic
Move to sternal	1-Smooth/methodical 2-Fighting mat, but controlled 3-Crashing, flopping over
Sternal phase	1-An organised pause 2-Non-existing 3-Multiple, with struggle
Moved to sternal	1-Methodical 2-An organised scramble 3-Used walls for support 4-Ricocheting off walls
Strength	1-Near full 2-Mildly rubbery 3-Dog sitting before standing 4-Repeated attempts due to weakness
Number of attempts to stand	Score = #
Balance and coordination	1-Solid 2-Moderate dancing 3-Reflex saves 4-Careening 5-Falls back down
Knuckling	1-None 2-Hindlimbs only 3-All four
Based on total score	1-Smooth, calm
	2-Calm but with some weakness, mild ataxia less than 10 minutes in duration
	3-As 2 but ataxia more than 10 minutes in duration
	4-Uncoordinated, with difficulties and ataxia, more than 3 attempts to stand
	5-Difficult recovery, unable to stand in the first 5 attempts; ataxia that can last for more than 20 minutes

Eight hundred and twenty-three horses were included in the study; 473 were geldings, 301 mares and 49 stallions. The age range was 1–33 years, the median age was 10 and the interquartile range (IQR) was 8 years; the weight range was 53–892 kg, the median was 526 and the IQR was 144 kg. Several breeds were included in the study, with Thoroughbred being the most represented one (158). ASA I category included 284, ASA II 285, ASA III 157, ASA IV 88 and ASA V 9 horses. Five hundred and eleven horses underwent elective procedures, whereas 312 underwent emergency surgery, with exploratory laparotomy for colic surgery being the most represented procedure.

The Spearman's correlation coefficient between the two systems was 0.88 (confidence interval 0.86 to 0.89; $P < 0.01$); overall k -weighted was 0.65. When horses were divided into groups based on ASA category, k -weighted was 0.64 for ASA I, 0.69 for ASA II, 0.59 for ASA III, 0.63 for ASA IV and 0.82 for ASA V. When horses were divided into groups based on type of surgery, k -weighted was 0.68 for elective and 0.59 for emergency procedures.

In 27 per cent of cases, D score was higher than Y, whereas only in 2 per cent of cases Y score was higher than D. In 528 (64 per cent) horses, the two scoring

systems gave the same score; in 282 (34 per cent) horses the score difference was only one point and in 13 (1.6 per cent) cases the difference was two points. Among these 13 cases, in 11 cases D and Y systems scored 5 and 3, respectively, and in two cases D system scored 4 and Y system scored 2. Most commonly, the disagreement between the two systems was for recovery scores 1–2 and 2–3.

Both Young and Taylor's² and Donaldson and others'³ systems have been used to score recoveries from general anaesthesia in clinical studies in horses, and they have been considered reliable⁴ and repeatable.⁷ Strong correlation between the two systems has already been reported⁸ and has been confirmed with our results; however, correlation does not reflect agreement, and therefore in our study k -weighted was calculated too. Substantial agreement ($k=0.65$) was reported between the two systems overall, but almost perfect agreement ($k=0.82$) was found only for horses belonging to the ASA V category, confirming what has been previously reported by Suthers and others.⁷

In our study, D system gave higher scores than Y system in 27 per cent of cases; this fact must be considered if multicentre studies are performed including facilities that use different RSS.

Clark-Price and others⁹ reported that subjective composite scales can lead different raters to different conclusions about the recovery status of the horse; ideally, an objective method should be developed in order to avoid inter-rater and intrarater variability, allowing comparison of studies performed at different centres.

The main limitation of our study is that people with different levels of experience were involved in scoring the recoveries. In fact, in our teaching hospital, the recovery score is often performed by students under the supervision of experienced equine anaesthetists (diplomates of the European or American College of Veterinary Anaesthesia and Analgesia) or anaesthesia residents, but the name of the person effectively scoring the recovery was not reported in the anaesthetic record; it was therefore not possible for the authors to record the level of experience of the observer or to include it in the analysis. However, there is evidence that the observer's experience does not affect recovery quality scoring.^{4 10}

Conclusions

A very strong correlation between Y and D recovery scoring systems exists. Their agreement was overall substantial, but the D system will generally give higher scores than the Y system. This fact should be considered when comparing studies using these two systems to evaluate recovery quality or if a multicentre study is performed.

Competing interests None declared.

Ethics approval Institutional ethical approval was obtained (VREC377).

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

© British Veterinary Association (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

References

- 1 JOHNSTON GM, EASTMENT JK, WOOD J, *et al*. The confidential enquiry into perioperative equine fatalities (CEPEF): mortality results of Phases 1 and 2. *Vet Anaesth Analg* 2002;29:159–70.
- 2 YOUNG SS, TAYLOR PM. Factors influencing the outcome of equine anaesthesia: a review of 1,314 cases. *Equine Vet J* 1993;25:147–51.
- 3 DONALDSON LL, DUNLOP GS, HOLLAND MS, *et al*. The recovery of horses from inhalant anaesthesia: a comparison of halothane and isoflurane. *Vet Surg* 2000;29:92–101.
- 4 VETTORATO E, CHASE-TOPPING ME, CLUTTON RE. A comparison of four systems for scoring recovery quality after general anaesthesia in horses. *Equine Vet J* 2010;42:400–6.
- 5 VALVERDE A, GUNKELT C, DOHERTY TJ, *et al*. Effect of a constant rate infusion of lidocaine on the quality of recovery from sevoflurane or isoflurane general anaesthesia in horses. *Equine Vet J* 2005;37:559–64.
- 6 SAKLAD M. Grading of patients for surgical procedures. *Anesthesiol* 1941;2:281–4.
- 7 SUTHERS JM, CHRISTLEY RM, CLUTTON RE. Quantitative and qualitative comparison of three scoring systems for assessing recovery quality after general anaesthesia in horses. *Vet Anaesth Analg* 2011;38:352–62.
- 8 LEECE EA, CORLETT F, BREARLEY JC. A comparison of recovery times and characteristics with sevoflurane and isoflurane anaesthesia in horses undergoing magnetic resonance imaging. *Vet Anaesth Analg* 2008;35:383–91.
- 9 CLARK-PRICE SC, LASCOLA KM, CARTER JE, *et al*. Assessment of agreement among diplomates of the American College of Veterinary Anesthesia and Analgesia for scoring the recovery of horses from anesthesia by use of subjective grading scales and development of a system for evaluation of the recovery of horses from anesthesia by use of accelerometry. *Am J Vet Res* 2017;78:668–76.
- 10 FARMER E JL. Perspective of recovery 'quality' in horses after general anaesthesia: Scottish Agricultural College and The University of Edinburgh, 2009. Bsc (Hons) Dissertation.



Retrospective evaluation of correlation and agreement between two recovery scoring systems in horses

Stefania Scarabelli and Eva Rioja

Veterinary Record 2018 182: 169 originally published online December 1, 2017

doi: 10.1136/vr.104546

Updated information and services can be found at:

<http://veterinaryrecord.bmj.com/content/182/6/169>

These include:

References

This article cites 9 articles, 0 of which you can access for free at:
<http://veterinaryrecord.bmj.com/content/182/6/169#ref-list-1>

Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See:
<http://creativecommons.org/licenses/by/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Anaesthesia](#) (2)
[Open access](#) (129)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>